

Absorption spectral structure in highly charged zirconium plasmas in water window soft x-ray spectral region




Takanori Miyazaki^{1,2}, Goki Arai², Hiroyuki Hara², Thanh-Hung Dinh², Takeshi Higashiguchi², Akinobu Irie², Chihiro Suzuki³, Daiji Kato³, Akira Sasaki⁴, Padraig Dunne¹, and Gerry O'Sullivan¹

¹University College Dublin, ²Utsunomiya University, ³National Institute for Fusion Science, ⁴Quantum Beam Science Center, Japan Atomic Energy Agency

Contact e-mail: takanori.miyazaki.cst@gmail.com


1. Introduction

We have investigated the absorption spectral structure of laser-produced highly charged zirconium plasmas in the water window soft x-ray spectral region. Structure due to absorption, which originates from opacity effects in the recombination phase, was observed at a delay time of around 100 ns.




2. Spectral structures of Zr plasmas

A. Single pulse LPP experiment




- (a) Experimental setup.
- (b) Zr spectra by single pulse Nd:YAG laser.
- (c) ZrO₂ spectra by single pulse CO₂ laser.

B. LHD experiment




C. HULLAC calculation




3. Spectral difference by dual lasers


A. Spectra



B. Difference spectral intensities to the single pulse



C. Water window soft x-ray source size



4. Summary

We have characterized the absorption spectral structure of dual laser-produced highly charged zirconium plasmas in water window soft x-ray spectral region due to the opacity effects in the recombination phase. Emissivity and opacity have been also evaluated in Zr plasmas.

Acknowledgements

A part of this work was performed under "Project for Bio-Imaging and Sensing at Utsunomiya University" from MEXT. One of the authors (T.H.) also acknowledges support from The Canon Foundation, Research Grant (Basic Research) on TEPCO Memorial Foundation. The UCD group was supported by Science Foundation Ireland International Co-operation Strategic Award 13/ISCA/2846.

